Technical support: order@acebiolab.com

Phone: 886-3-2870051

Ver.1 Date: 20191114

IL17A Polyclonal Antibody


Cat# A340753- 20/50/100 ug

Storage at -20 °C


INFORMATION

Product Name Cat. No. A340753 Product type Primary antibodies Species reactivity Human, Mouse, Rat Clonality polyclonal Host Rabbit Size 20 ug/50 ug/100 ug Uniprot Q16552 Immunogen Recombinant protein of human IL17A Concentration 1 mg/ml
Product type Primary antibodies Species reactivity Human, Mouse, Rat Clonality polyclonal Host Rabbit Size 20 ug/50 ug/100 ug Uniprot Q16552 Immunogen Recombinant protein of human IL17A Concentration 1 mg/ml
Species reactivity Human, Mouse, Rat Clonality polyclonal Host Rabbit Size 20 ug/50 ug/100 ug Uniprot Q16552 Immunogen Recombinant protein of human IL17A Concentration 1 mg/ml
Clonality polyclonal Host Rabbit Size 20 ug/50 ug/100 ug Uniprot Q16552 Immunogen Recombinant protein of human IL17A Concentration 1 mg/ml
Host Rabbit Size 20 ug/50 ug/100 ug Uniprot Q16552 Immunogen Recombinant protein of human IL17A Concentration 1 mg/ml
Size 20 ug/50 ug/100 ug Uniprot Q16552 Immunogen Recombinant protein of human IL17A Concentration 1 mg/ml
Uniprot Q16552 Immunogen Recombinant protein of human IL17A Concentration 1 mg/ml
Immunogen Recombinant protein of human IL17A Concentration 1 mg/ml
Concentration 1 mg/ml
Tested applications WB, IHC
Application WB: 1:500-1:2000, IHC: 1:50-1:200
Conjugation None
Purification Affinity purified
Molecular Weight Calculated MW: 17 kDa, Observed MW: 37 kDa
Stock buffer PBS with 0.02% sodium azide and 50% glycerol pH 7.4.
Storage instruction Store at -20°C. Avoid freeze / thaw cycles.
Alias CTLA-8; CTLA8; IL-17; IL-17A; IL17
Expression Restricted to activated memory T-cells.
The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This
cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine
Background can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the
production of nitric oxide (NO). High levels of this cytokine are associated with several chronic
inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.






Western blot analysis of extracts of various cell lines with IL17A Polyclonal Antibody.



Image

Immunohistochemistry of paraffinembedded mouse kidney with IL17A Polyclonal Antibody.



Immunohistochemistry of paraffinembedded human stomach with IL17A Polyclonal Antibody.

PRODUCT USE LIMITATION

These products are intended for research use only.

