



# Rabbit Anti-CD16 antibody

Cat# A18514BS


Upon receipt, store at -20°C . Avoid repeated freeze.

## INFORMATION


<b>Product Name</b>	Rabbit Anti-CD16 antibody
<b>Cat. No.</b>	A18514BS
<b>Size</b>	20 µl ,100 µl
<b>Product type</b>	Primary Antibody
<b>Species Reactivity</b>	Hu, Ms, Rb
<b>Immunogen Range</b>	145-195/254
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Tested applications</b>	Western Blot: 1/300 - 1/1000. IHC-p: 1:200-1:400. FCM: 1:20-1:100. IF: 1/50-1/200
<b>Conjugation</b>	Non-Conjugated
<b>Purification</b>	Purified by Protein A.
<b>Concentration</b>	1 mg/mL
<b>Storage buffer</b>	Aqueous buffered solution containing 1% BSA, 50% glycerol and 0.09% sodium azide.
<b>Alias</b>	CD16; FCG3; CD16A; FCGR3; IGFR3; IMD2; FCR-1; FCRIII; FCGRIII; FCRIIIA; Low affinity immunoglobulin gamma Fc region receptor III-A; CD16a antigen; Fc-gamma RIII-alpha; Fc-gamma RIII; Fc-gamma RIIIa; IgG Fc receptor III-2; FCGR3A



Lane 1: Lung lysates; Lane 2: Spleen lysates probed with CD16 Polyclonal Antibody, Unconjugated at 1:300 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at 1:20000 for 60 min at 37°C.




Formalin-fixed and paraffin embedded mouse spleen labeled with Rabbit Anti IGFR3/CD16 Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining.



Formalin-fixed and paraffin embedded human glioma tissue labeled with Anti IGFR3/CD16 Polyclonal Antibody, Unconjugated (bs-6028R) at 1:200 followed by conjugation to the secondary antibody and DAB staining.

Image



Formalin-fixed and paraffin-embedded human lung cancer labeled with Rabbit Anti-CD16 Polyclonal Antibody, Unconjugated 1:200, overnight at 4°C, The secondary antibody was Goat Anti-Rabbit IgG, Cy3 conjugated used at 1:200 dilution for 40 minutes at 37°C.

### **PRODUCT USE LIMITATION**

These products are intended for research use only.