Technical support: order@acebiolab.com

Phone: 886-3-2870051

Ver.1 Date: 20180222

## **ER Stress Antibody Sampler Kit**

Cat# AK0155

Upon receipt, store at -20°C. Avoid freeze/thaw cycles.

## **PRODUCT DESCRIPTION**

Secretory and transmembrane proteins are synthesized on polysomes and translocate into the endoplasmic reticulum (ER) where they are often modified by the formation of disulfide bonds, amino-linked glycosylation and folding. The ER contains a pool of molecular chaperone proteins including calnexin, BiP and protein disulfide isomerase (PDI). Calnexin is an ER membrane, calcium-binding protein that retains newly synthesized glycoproteins inside the ER to ensure proper folding and quality control. Irregular protein folding within the ER increases BiP synthesis, which binds misfolded proteins to prevent them from forming aggregates and to assist them to refold properly. PDI catalyzes the formation and isomerization of disulfide bonds required for a protein to reach its native state. Studies have found that the resident ER protein endoplasmic oxidoreductin-1 (Ero1) provides oxidizing potential to the ER in Saccharomyces cerevisiae. Ero1-L lpha is an ER membrane-associated N-glycoprotein that promotes oxidative protein folding. Disruptions of ER homeostasis leads to the accumulation of unfolded proteins. The ER has developed an adaptive mechanism called the unfolded protein response (UPR) to counteract compromised protein folding. This is regulated by proteins such as the membrane-bound transcription factor protease site 2 (MBTPS2) and the serine/threonine kinase IRE1. The PERK eIF2  $\alpha$  kinase is an ER resident transmembrane protein that couples ER stress signals to translation inhibition. ER stress increases PERK activity, which phosphorylates eIF2 lpha to reduce protein translation. PERK activation during ER stress correlates with autophosphorylation of its cytoplasmic kinase domain. Phosphorylation of PERK at Thr980 can serve as a marker for its activation status. During ER stress, the level of CHOP expression is elevated and CHOP functions to mediate programmed cell death.

## **PRODUCT INCLUDES**

Cat No.	Product name	Quantity	Applications	Reactivity	Host
A340552	HSPA5 Polyclonal Antibody	20μL	WB, IHC, IF, ELISA	Human, Mouse, Rat, Monkey	Rabbit
A340454	CANX Polyclonal Antibody	20μL	WB, IHC, IF, ELISA	Human, Mouse, Rat	Rabbit
A340482	CHOP Polyclonal Antibody	20μL	WB, IHC, IF, ELISA	Human, Mouse	Rabbit



A340731	PERK Polyclonal Antibody	20μL	WB, IHC-p, ELISA	Human, Mouse, Rat	Rabbit
A1013s	Goat Anti-Rabbit IgG (H+L) (peroxidase/HRP conjugated)	120μL	WB, ELISA	Rabbit	Goat

## **PRODUCT USE LIMITATION**

These products are intended for research use only.

