Technical support: order@acebiolab.com

Phone: 886-3-2870051

Ver.1 Date: 20180222

PDGF Receptor α Antibody Sampler Kit

Cat# AK0218

Upon receipt, store at -20°C. Avoid freeze/thaw cycles.

PRODUCT DESCRIPTION

Platelet derived growth factor (PDGF) family proteins exist as several disulphide-bonded, dimeric isoforms (PDGF AA, PDGF AB, PDGF BB, PDGF CC, and PDGF DD) that bind in a specific pattern to two closely related receptor tyrosine kinases, PDGF receptor α (PDGFR α) and PDGF receptor β (PDGFR β). PDGFR α and PDGFR β share 75% to 85% sequence homology between their two intracellular kinase domains, while the kinase insert and carboxy-terminal tail regions display a lower level (27% to 28%) of homology. PDGFR lpha homodimers bind all PDGF isoforms except those containing PDGF D. PDGFR eta homodimers bind PDGF BB and DD isoforms, as well as the PDGF AB heterodimer. The heteromeric PDGF receptor α/β binds PDGF B, C, and D homodimers, as well as the PDGF AB heterodimer. PDGFR lpha and PDGFR eta can each form heterodimers with EGFR, which is also activated by PDGF. Various cells differ in the total number of receptors present and in the receptor subunit composition, which may account for responsive differences among cell types to PDGF binding. Ligand binding induces receptor dimerization and autophosphorylation, followed by binding and activation of cytoplasmic SH2 domain-containing signal transduction molecules, such as GRB2, Src, GAP, PI3 kinase, PLC γ , and NCK. A number of different signaling pathways are initiated by activated PDGF receptors and lead to control of cell growth, actin reorganization, migration, and differentiation. Tyr751 in the kinase-insert region of PDGFR β is the docking site for PI3 kinase. Phosphorylated pentapeptides derived from Tyr751 of PDGFR β (pTyr751-Val-Pro-Met-Leu) inhibit the association of the carboxy-terminal SH2 domain of the p85 subunit of PI3 kinase with PDGFR β . Tyr740 is also required for PDGFR β -mediated PI3 kinase activation.

PRODUCT INCLUDES

Cat No.	Product name	Quantity	Applications	Reactivity	Host
A340626	PDGFRA Polyclonal Antibody	20μL	WB, IHC, IF, ELISA	Human, Mouse, Rat	Rabbit
A340311	Phospho-PDGFRA (Tyr849) Polyclonal Antibody	20μL	WB, IHC, ELISA	Human, Mouse, Rat	Rabbit
A1013s	Goat Anti-Rabbit IgG (H+L) (peroxidase/HRP conjugated)	120μL	WB, ELISA	Rabbit	Goat



PRODUCT USE LIMITATION

These products are intended for research use only.

