Technical support: <a href="mailto:order@acebiolab.com">order@acebiolab.com</a>

Phone: 886-3-2870051

Ver.1 Date: 20180222

## IL-6Rα Polyclonal Antibody

Cat# A21668PI

Upon receipt, store at -20°C. Avoid repeated freeze.

## **INFORMATION**

Product Name	IL-6Rα Polyclonal Antibody		
Cat. No.	A21668PI		
Size	100ug/50ug/20ug		
Uniprot	P08887		
Product type	1 9999		
	Primart antibody		
Species Reactivity	Human  The anticorum was produced against synthesized portide derived from the Internal		
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal		
	region of human IL6R. AA range:221-270		
Host	Rabbit		
Concentration	1 mg/ml		
Clonality	Polyclonal		
Tested applications	WB,IHC-p,IF(paraffin section),ELISA		
Application	IF(paraffin section) 1:50-200 IHC-p: 100-300.Western Blot: 1/500 - 1/2000. ELISA:		
	1/10000. Not yet tested in other applications.		
Purification Method	The antibody was affinity-purified from rabbit antiserum by affinity-		
	chromatography using epitope-specific immunogen.		
Molecular Weight	53KD		
Storage instruction	-20°C/1 year		
Alias	IL6R; Interleukin-6 receptor subunit alpha; IL-6 receptor subunit alpha; IL-6R		
	subunit alpha; IL-6R-alpha; IL-6RA; IL-6R 1; Membrane glycoprotein 80; gp80;		
	CD126		
Image		Western Blot analysis of AD293, 4T1	
	AD293 4T1 138— 100— 70—	cells using IL-6Rα Polyclonal Antibody	
		Secondary antibody was diluted at	
	55 ILGR	1:20000	
	45		
	35		
	25		



Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3, Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3, Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3, Secondary antibody was diluted at 1:200(room temperature, 30min).		
1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at		·
overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at 1:200(was used for antigen retrieval).  3,Secondary antibody was diluted at d		paraffin-embedded Human spleen.
2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at		1, Antibody was diluted at 1:200(4°
pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at		overnight).
3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at	n) 1	2, High-pressure and temperature EDTA,
1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		pH8.0 was used for antigen retrieval.
Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		3,Secondary antibody was diluted at
paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		1:200(room temperature, 30min).
1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at		Immunohistochemical analysis of
overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		paraffin-embedded Human spleen.
2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at		1, Antibody was diluted at 1:200(4°
pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		overnight).
3,Secondary antibody was diluted at 1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at		2, High-pressure and temperature EDTA,
1:200(room temperature, 30min).  Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		pH8.0 was used for antigen retrieval.
Immunohistochemical analysis of paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		3,Secondary antibody was diluted at
paraffin-embedded Human spleen.  1, Antibody was diluted at 1:200(4° overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		1:200(room temperature, 30min).
1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at		Immunohistochemical analysis of
overnight).  2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		paraffin-embedded Human spleen.
2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		1, Antibody was diluted at 1:200(4°
pH8.0 was used for antigen retrieval.  3,Secondary antibody was diluted at		overnight).
3,Secondary antibody was diluted at		2, High-pressure and temperature EDTA,
		pH8.0 was used for antigen retrieval.
1:200(room temperature, 30min).		3,Secondary antibody was diluted at
	Y to the second	1:200(room temperature, 30min).

## **PRODUCT USE LIMITATION**

These products are intended for research use only.

